

New herbage plant cultivar

B. Legumes

8. Lucerne

(a) *Medicago sativa* L. (lucerne) cv. Sequel HR

Reg. No. B-8a-24. Registered on 10 November, 1997.

Originators: BRAY, R.A.¹ and IRWIN, J.A.G.²

¹CSIRO Tropical Agriculture, 306 Carmody Rd, St Lucia, Qld 4067, Australia.

²Botany Department, The University of Queensland, Brisbane, Qld 4069, Australia.

Registrar: Kelman, W.M.

CSIRO Plant Industry, GPO Box 1600, Canberra, ACT 2601, Australia.

Released by CSIRO Tropical Agriculture and The University of Queensland.

Published in the Australian Journal of Experimental Agriculture 1998, **37**, 000–000.

Origin

Sequel HR (“Highly Resistant”) is a highly winter-active lucerne derived from the cultivar Sequel by selection for high levels of resistance to anthracnose (*Colletotrichum trifolii* Bain. et Essary). Seedlings of a population known as SS2 (derived by selection from Sequel for resistance to *Stemphylium vesicarium* Simmonds) were selected under controlled conditions for resistance to *C. trifolii* (seedling inoculation) and *Phytophthora medicaginis* Hansen et Maxwell (stem inoculation; Irwin *et al.* 1980). Approximately 100 seedlings that met the selection criteria were selected and intercrossed under glasshouse conditions. The progeny were subjected to further selection for the two diseases on a maternal line basis (one selection from each line where possible) and 80 selections were intercrossed to form the population tested as CS93-1. This provided the breeders seed for Sequel HR. CSIRO Tropical Agriculture will maintain breeders seed. Release of Sequel HR was endorsed by the Queensland Herbage Plant Liaison Committee in August 1996. Sequel HR has provisional protection under Plant Breeders Rights legislation (Application No 95/142).

Morphological description

Sequel HR is a highly winter-active cultivar (dormancy group 9); it is similar in appearance to Sequel (Oram 1990) and taller than Aquarius, CUF 101 and Sceptre (all dormancy group 9). Flowers are purple and time to commencement of flowering in spring is similar to that for Sequel and slightly earlier than for Aquarius, CUF 101 and Sceptre. Seed size of Sequel HR (approximately 430 000 per kg) is similar to that for Sceptre and Aquarius and smaller than that for Sequel and CUF 101.

Agronomic characters

Sequel HR was developed as a replacement for Sequel in the Queensland subtropics. Sequel HR has a very high level of anthracnose resistance (70–80% resistant plants), significantly greater than that of Sequel, Aquarius, Sceptre, CUF 101 and Trifecta, but similar to that of Pioneer L69. Sequel HR is resistant to *Phytophthora* (40% resistant plants) with a level similar to that of Sequel and Sceptre but not as high as that of Aquarius.

Sequel HR was included in three trials conducted by Dr Rex Williams of NSW Agriculture, Yanco, to evaluate resistance to spotted alfalfa aphid (*Therioaphis trifolii* (Monell) f. *maculata*). Sequel HR was rated as resistant in the three tests, with a mean of 35% resistant plants compared with 40% for Sceptre, 30% for Aquarius and 5% for Hunter River (R. W. Williams, personal communication).

Having been derived from Sequel by selection while maintaining a broad genetic base, Sequel HR is expected to retain the broad adaptation to Australian lucerne growing areas exhibited by Sequel. Like Sequel, Sequel HR should be well suited to a range of situations from hay production through to use as a ley legume in cereal rotations. It was one of the highest yielding

cultivars in an irrigated trial conducted at Gatton Research Station by Mr K.F. Lowe on ground known to be heavily infested with *Phytophthora medicaginis*. Over the three years of the trial (1993–1996), first generation Sequel HR gave yields similar to those of Sequel, Aquarius and Sceptre and outyielded Trifecta by 12% and Hunter River by 46%. In another trial at Gatton Research Station established in 1995, Sequel HR was also among the group of highest yielding cultivars (K.F. Lowe, personal communication). Due to drought conditions, anthracnose was not a problem in these trials. In more normal years, the extra anthracnose resistance possessed by Sequel HR over Sequel should provide a yield advantage.

Acknowledgements

Technical assistance was provided by L. Hart and J. Mackie. Data from the Gatton trial were supplied by K.F. Lowe, Queensland Department of Primary Industries and data on resistance to spotted alfalfa aphid were provided by Dr R.W. Williams of NSW Agriculture.

References

- IRWIN, J.A.G., MAXWELL, D.P. and SMITH, R.R. (1980) Correlation between root and stem reaction of alfalfa to *Phytophthora megasperma* f. sp. *medicaginis*. *Phytopathology*, **70**, 987–990.
- ORAM, R.N. (1990) Sequel lucerne. In: Oram, R.N. (ed.) *Register of Australian Herbage Plant Cultivars*. 3rd Edn. pp. 204–205. (CSIRO: Melbourne).