

New herbage plant cultivars

A. Grasses

23. Digitaria

(b) *Digitaria milanjiana* (Rendle) Stapf. (finger grass) cv. Jarra

Reg. No. A-23b-1. Registered on January 15, 1993.

Originators: HALL, T.J.,¹ WALDUCK, G.D.² and WALKER, R.W.³

¹ Queensland Department of Primary Industries, PO Box 308, Roma, Qld 4455, Australia.

² PO Box 943, Innisfail, Qld 4860, Australia.

³ Queensland Department of Primary Industries, PO Box 1054, Mareeba, Qld 4880, Australia.

Registrar: Oram, R.N.

CSIRO Division of Plant Industry, GPO Box 1600, Canberra, ACT 2601, Australia.

Released by Queensland Department of Primary Industries.

Published in the *Australian Journal of Experimental Agriculture*, 1993, 33, 000.

Origin

Jarra (*Digitaria milanjiana* CPI 59745) was collected by Mr R.W. Strickland of CSIRO (No. RWS 465) on February 12, 1972 in Malawi, 46 km west of Blantyre on the Chikwawa road (latitude 16°S, altitude 61 m, rainfall 700 mm) on a freely drained loam soil (pH 7.5) with a 10–20 cm surface horizon (Anon. 1973), in a run-on grassland area of *Urochloa*, *Brachiaria*, *Rhynchosia*, *Tephrosia* and *Alysicarpus* species (R.W. Strickland, personal communication).

Digitaria milanjiana (woolly finger or milanje finger grass in Africa) is a variable, tropical perennial species (Hacker 1983), with rhizomatous and stoloniferous or densely tufted ecotypes (Chippindale 1955). It is distributed in tropical eastern Africa, from Kenya to South Africa and in Angola, at altitudes from 0–2070 m. The climatic range is from semi-arid tropics (450 mm rainfall) to wet equatorial (1700 mm). It commonly grows in grasslands in moderate rainfall areas on sandy loam. It also occurs naturally in open woodlands on black heavy soils

as well as on sandy soils, and can be a weed in cultivations. It is readily accepted by cattle (Ibrahim and Kabuye 1987). In trials, *D. milanjiana* has had some success in pure stands (Roberts 1970) and in legume mixtures with *Trifolium repens* and *Lotononis bainesii* (Clatworthy 1970).

Stolons of CPI 59745 were transplanted from CSIRO Lansdown Research Station to the QDPI nursery at Walkamin Research Station in 1982. Stolon material from this nursery was planted in a grass adaptation experiment in a Verano and Seca stylo pasture at Springs Road near Mareeba in 1983. CPI 59745 persisted over five years and spread under continuous grazing for the last three years.

Seed production of the *Digitaria* collection was assessed in swards at Walkamin and grasses in the Walkamin nursery were tested for resistance to the burrowing nematode (*Radopholus similis* (Cobb) Thorne), a serious economic pest of bananas (Broadley 1977). As *D. milanjiana* was found resistant to the nematode (W. O'Donnell, personal communication), the ten most successful accessions from the grazed adaptation and seed production experiments were evaluated at South Johnstone Research Station for growth rate, stolon development, spreading ability and capacity to compete with weeds (mainly *Mimosa pudica*, sensitive weed) in a banana plantation. CPI 59745 and two other stoloniferous lines were selected from this high rainfall site to be evaluated in commercial banana crop rotations at Jarra Creek near Tully, the Palmerston district near Innisfail and near Mission Beach. CPI 59745 was the most strongly stoloniferous, vigorous and competitive of the three accessions, and release by the Queensland Department of Primary Industries as cultivar Jarra was endorsed by the Queensland Herbage Plant Liaison Committee in August 1991. Planting material has been distributed from Walkamin, South Johnstone and Jarra Creek.

Morphological description

Jarra is a stoloniferous perennial with long, mauvish-coloured stolons 10–15 cm between culms. Basal cataphylls with woolly hairs. Flowering culms erect, tufted, 50–120 cm tall, unbranched, terminated by a solitary inflorescence, 2–4 noded. Nodes glabrous, forming a purple-coloured ring. Internodes spongy, glabrous, longer than associated leaf sheaths at culm base or shorter than these towards culm apex. Leaf sheaths keeled, pilose, with tubercle-based hairs, with finely scaberulous nerves. Ligule a membrane, 2–2.5 mm long, slightly erose, acute. Collar glabrous. Leaf blades flat, 50–200 mm x 3–8 mm, linear, pilose with tubercle-based hairs, tapering to a narrow apex, attenuate and ciliate at the base, with scabrous margins. Inflorescence of subdigitate racemes. Main axis 15–20 cm long, scabrous. Primary branches with pubescent axils, spreading, with spikelets appressed to the rachis and neatly arranged, 60–130 mm long, 1–2 mm wide, triquetrous and winged, scabrous on the margins. Pedicels 0.2–0.7 mm long, distinctly angled, scabrous. Pedicel apices glabrous, concave, truncate. Disarticulation at the base of the spikelet. Spikelets green to pale brown or purple, abaxial, densely overlapping, paired, with short and long pedicels, 120–180 on a typical primary branch, dorsiventrally compressed, attenuate at base, lanceolate, 2.6–3 x 1 mm. Lower glume 0.2–0.3 mm long, deltoid, nerveless, membranous, smooth, glabrous, acute. Upper glume 0.7–1.2 mm long, lanceolate, 3-nerved, membranous, glabrous or villous on nerves, acute. Lower floret neuter: lemma 2.8–3.2 x 1 mm, lanceolate, membranous, with the first internode space wider than the second, 5-nerved, the nerves scabrous, with involute margins, glabrous or villous on nerves, smooth or very finely scabrous, acute; palea *ca.* 0.2 mm long, vestigial. Upper floret perfect, slightly shorter than the lower floret; lemma shiny, 2.3–2.4 mm long, green to brown, membranous, smooth, elliptic, shiny, glabrous, acute; palea membranous, smooth, enclosed at its apex by the lemma. Stamens 3 with anthers *ca.* 1.5 mm long and purple. Caryopsis *ca.* 2 mm long, pale brown, ellipsoid (B.K. Simon, personal communication).

In Africa, *D. milanijana* has been described by Henrard (1950); Chippindale (1955); Bogdan

(1977); Clayton and Renvois (1982); and Ibrahim and Kabuye (1987). Synonyms for *D. milanijana* include: *D. setivalva*, (Stent), *D. boivinii* (Henr.), *D. endlichii* (Mez.), and *D. mombasana* (C.E. Hubbard). Introductions to Australia are highly variable, differing in lemma setigerousness, growth habit, and leaf and stem hairiness (Hacker 1983). *D. milanijana* is distinguished from the *D. eriantha* group in having scabrid nerves on the lower lemma and is largely tropical; *D. eriantha* is subtropical (Clayton and Renvois 1982).

Jarra can be distinguished from other *D. milanijana* accessions (e.g. CPI 59755), *D. swynertonii* CPI 59749, and *D. smutsii* cv. Premier, by electrophoretic analysis of bulked seed lots (5 seeds) using the SDS-PAGE method (Vieritz 1990). *D. milanijana* is a diploid with $2n = 18$ chromosomes.

Agronomic characters

In a grazed stylo pasture at Mareeba, Jarra has been among the better adapted *D. milanijana* accessions between 1983 and 1987 (T.J. Hall and R.W. Walker, unpublished data). Native grasses are not persistent in these sown legume pastures in the dry tropics of northern Australia and grazing-tolerant grasses are required to withstand increased stocking pressure. Strongly stoloniferous species, such as Jarra, may be useful in this role.

Jarra has persisted and spread on a range of soil types from coarse granitic sandy loams, alluvial loams and shallow solodics, to a eucrozem. Growth has been greatest on the more fertile soils and at higher rainfall. The species is reported to be intolerant of waterlogging. On a loam soil at Tully, spread and vigour of Jarra were poorest on the lower slope of a seasonally-wet creek levee where sedges (*Cyperus* spp.) were prolific. Growth was rapid higher up the slope where soil internal drainage was good.

The banana cropping system on the wet coast of north Queensland requires a species to rotate with bananas that will break the burrowing nematode (*R. similis*) cycle, while maintaining weed control and erosion protection. *D. milanijana* was found to satisfy these three criteria.

Within three months of planting stolons of Jarra at Tully on a cultivated soil with little weed competition, plants established and produced

multi-branched stolons 2.6 m long with new plants (mean height 31 cm) established at the first ten nodes. The spacing of these new plants averaged 12 cm (commonly 7–17 cm) apart. There were commonly 18–32 tillers per original plant, although some had over 80. These tillers readily developed into rooted stolons as they elongated on bare soil. There was 100% soil cover within six months, and within 12 months, Jarra yielded 10.4 t/ha DM, with 68% green material. This green leaf had 2.11% N, 0.28% P, 0.16% S and 0.84% Ca in June. In the second year, dry matter yield was 15.9 t/ha DM. There were scattered *Cassia obtusifolia* (sickle pod) bushes growing above the 1 m tall grass, but no herbaceous weeds grew under the dense canopy. There were no *R. similis* present.

Seed production of *D. milanjiana* is not well recorded. However, germination improves with over five months storage. The removal of the palea and lemma improves germination and dehulled seeds can germinate soon after harvest. Gibberellin stimulates germination of dehulled but not intact seeds (Baskin *et al.* 1969). Poor seed production of the stoloniferous accessions has been one limitation to more widespread testing. Jarra has produced commercially acceptable yields (about 100 kg/ha) of good quality, clean seed by header-harvesting at Tully and Walkamin. A pure seed sample contains about 2,000,000 seeds/kg. It is not yet known if Jarra propagated by seed would remain true to type.

Vegetative establishment provides rapid soil cover after removing bananas. Jarra stolons planted in summer in a strip between the old banana rows will give complete cover in three months. A grass rotation lasting two years with no weeds which are hosts to the burrowing nematodes will break the nematode cycle, a much cheaper alternative to chemical controls. Without nematode control, yield losses occur in ratoon crops due to plants falling over and a reduced bunch size (Broadley 1977). Alternative pasture species such as signal (*Brachiaria decumbens*) and guinea (*Panicum maximum*) grasses are hosts to the burrowing nematode. Herbicides (glyphosate and Gramoxone) can be used to kill Jarra to

prepare for planting the following banana crop.

Jarra is the first Australian cultivar of the species and is proposed for use as a pasture ley in banana cropping areas of Queensland and as a pasture plant in favourable environments of the seasonally-dry tropics and wet coastal tropics.

Acknowledgements

We thank Dr J.M. Hopkinson for seeding evaluation, Mr L. Schipke and Mr W. O'Donnell for nematode testing, and Messrs G. Flegler, H. Sly and C. Rick for use of land. Mr G. Flegler developed the commercial establishment methods. The Australian Meat and Livestock Research and Development Corporation provided funds for the grass selection experiment.

References

- ANON (1973) *Australian Plant Introduction Review* 9: 70 A. (CSIRO Division of Plant Industry: Canberra).
- BASKIN, J.M., SCHANK, S.C. and WEST, S.H. (1969) Seed dormancy in two species of *Digitaria* from Africa. *Crop Science* 9: 584–586.
- BOGDAN, A.V. (1977) *Tropical Pasture and Fodder Plants (Grasses and Legumes)*. (Longman: London).
- BROADLEY, R.A. (1977) *Nematode control in bananas and tobacco grown in north Queensland*. Ph.D Thesis, University of London.
- CHIPPINDALE, L.K.A. (1955) A guide to the identification of grasses in South Africa. In Meredith, D. (ed.) *The Grasses and Pastures of South Africa*. (South Africa Central News Agency).
- CLATWORTHY, J.N. (1970) A comparison of legume and fertilizer nitrogen in Rhodesia. *Proceedings of the XI International Grassland Congress*, Surfers Paradise. pp. 408–411.
- CLAYTON, W.D. and RENVOISE, S.A. (1982) *Flora of Tropical East Africa*. Part 3. Gramineae. (Crown Agents Overseas Govts Admin.: London).
- HACKER, J.B. (1983) Inheritance of stolon development, rhizome development and setigerous lemmas in the *Digitaria milanjiana* complex, and its taxonomic significance. *Australian Journal of Botany*, 31: 357–369.
- HENRARD, J.T. (1950) Monograph of the genus *Digitaria*. University pers Leiden, Leiden, pp. 456–461 and 962.
- IBRAHIM, K.M. and KABUYE, C.H.S. (1987) *An Illustrated Manual of Kenya Grasses*. pp. 127 and 413. (FAO, United Nations: Rome).
- ROBERTS, O.T. (1970) A review of pasture species in Fiji. I. Grasses. *Tropical Grasslands*, 4: 129–137.
- VIERITZ, A.M. (1990) Electrophoretic analysis of *Digitaria milanjiana* and *D. smutsii* by SDS-PAGE. *Seed Research Project Report 9.14*, Queensland Department of Primary Industries, Brisbane.